

# Masodine® Concentrate

Teat dip / spray and udder wash solution 2.15% w/v



Economical concentrated formulation, helps to reduce bactericidal contamination. Effective aid in the control of mastitis in dairy herds.

UK Authorised Veterinary Medicine.

Formulated to help reduce bacterial contamination that causes mastitis in dairy herds.

- Reduces the risk of infection by leaving an antiseptic film of Iodine adhering to teats which have been treated by dipping or spraying.
- Use as a post-milking dip or spray at full dilution rate of 1:3.
- May also be used as a pre-milking dip or spray diluted at a rate of 1:7.
- Contains 20% Glycerol and Sorbitol (5% when diluted) to help keep teats soft and supple and to aid in the prevention and healing of sores and cracks.

#### TEAT DIP OR SPRAY: (5,000 ppm minimum available Iodine)

Dilute 1:3 parts clean water and mix well.

**TEAT DIPPING:** Fill teat cup about two thirds full with the prepared solution. Dip teats of every cow immediately after each has been milked, making sure that the full length of the teat is immersed. Top up with fresh solution as required. Empty and wash out cups after milking.

**TEAT SPRAYING:** Immediately after milking, spray the entire surface of each teat of every cow with prepared solution.

#### UDDER WASH AND CLUSTER DIP: (40 ppm minimum available Iodine)

Dilute one part product with 500 parts clean water (20 ml per 10 litres or 1 fl oz in 3 gal). To prevent cross infection, individual cloths or disposable paper towels are strongly recommended. Dry teats before milking. Immerse clusters and agitate for at least half a minute. Rinse in clean water before use. Wash and dry udders and teats before milking.

#### ALTERNATIVE USAGE - PRE-MILKING DISINFECTION: (2,500 ppm minimum available Iodine)

Dilute 1:7 parts clean water and mix well. Ensure udder and teats are clean and dry before pre-dipping.

**PRE-MILKING TEAT DIPPING:** Fill teat dipping cup 2/3 full with the 1:7 solution. Dip teats of every cow making sure that the full length of the teat is immersed. Allow 15 seconds contact time, then using a single service paper towel, wipe and dry teat thoroughly, before continuing with normal milking routine. Top up with fresh solution as required. Empty and wash out cups after milking.

**PRE-MILKING TEAT SPRAYING:** Spray the entire surface of each teat of every cow with 1:7 solution. Allow 15 seconds contact time, then using a single service paper towel, wipe and dry teat thoroughly, before continuing with normal milking routine.

25 L

**Order Code: R071KEV**

Pack: 25 litre

200 L

**Order Code: R071NEV**

Pack: 200 litre

1000 L

**Order Code: R071SEV**

Pack: 1000 litre

#### PRODUCT INFORMATION:

Appearance:	Dark brown liquid
Odour:	Faint Iodine
pH - undiluted:	4.0
Shelf life:	2 years

#### QUALITY AND ENVIRONMENTAL ASSURANCE:

This product is manufactured in the U.K. by EVANS VANODINE INTERNATIONAL PLC under an ISO 9001 Quality Management System Cert. No. FM 09535 and an ISO 14001 Environmental Management System Cert. No. EMS 506072 registered by the British Standards Institution.

**REVISION DATE:** 31/07/25



Scan here to view safety data information

