

---

# MICROBIOLOGICAL PROFILE



**Evafoam**<sup>™</sup>

Ready-to-use pre/post-milking foaming teat disinfectant

Edition 1: October 2023

# EVAFOAM MICROBIOLOGICAL PROFILE

## INTRODUCTION

**EVAFOAM** is a ready-to-use pre/post-milking foaming teat disinfectant.

**EVAFOAM** is a biocidal product which may be used as a dip, spray or foam for the cleaning of cow's teats before and after milking.

**EVAFOAM** disinfects cows' teats and udder.

**EVAFOAM** passes EN 1656. tested against common mastitis pathogens in the presence of milk.

**EVAFOAM** is recommended for use as part of a Dairy Hygiene Programme.

Effective aid in the control of mastitis in dairy herds		Can be used for pre and post milking
Contains Allantoin and Glycerol to help smooth and soften the skin	Contains the biocides Chlorhexidine Gluconate and Lactic Acid	Tested in a UKAS accredited laboratory

## EVAFOAM - EFFICACY SUMMARY

EVAFOAM has been tested and proven to be effective against a range of micro-organisms. European Standard (EN\*) test methods were used to prove efficacy against bacteria.

The UKAS accredited Microbiology Laboratory at Evans Vanodine International plc. (Testing number 1108) performed tests with bacteria and yeast.

The following tables include information of relevant, applicable test methods, conditions, contact times and organisms.

\*EN - European Norm

Published in the UK as BS EN by the British Standards Institution.



# EVAFOAM MICROBIOLOGICAL PROFILE

## ACTIVITY AGAINST BACTERIA

BACTERIAL TEST PROFILE				
ORGANISMS	TEST METHOD	TEMP (°C)	CONTACT TIME (MINUTES)	SOILING LEVEL
<i>Escherichia coli</i>	EN 1656	30	30 seconds	Pre milking
<i>Staphylococcus aureus</i>				
<i>Streptococcus uberis</i>				
<i>Escherichia coli</i>			5	Post milking
<i>Staphylococcus aureus</i>				
<i>Streptococcus uberis</i>				

# EVAFOAM MICROBIOLOGICAL PROFILE

## VETERINARY DISINFECTANT TEST METHODS

Veterinary disinfectants can be used in a variety of areas e.g. the breeding, husbandry, production, transport and disposal of all animals except when in the food chain following death and entry to the processing industry.

There are two types of laboratory test methods for livestock disinfectants, suspension and surface methods. As a minimum for general hygiene purposes, products should be effective against bacteria and yeast.

The scope of veterinary EN test methods does not specify application of the product but does include disinfection by immersion and surface disinfection by wiping, spraying, foaming or other means. It does not include aerial disinfection.

The interfering substances used in EN test methods are described as low or high level soiling for disinfectants and as pre and post milking for teat disinfectants in the veterinary test methods. They simulate levels of soiling encountered in practical, real-life situations.

## EN TEST METHODS

TEST REFERENCE		TEST TYPE	ORGANISM	TEST PASS CRITERIA
EN 1656	For bactericidal activity.	Suspension	Bacteria	≥5 log reduction

# EVAFOAM MICROBIOLOGICAL PROFILE

## LOG REDUCTION

Products claiming they will kill 99.9% of bacteria sounds extremely efficient, however it does not prove that a product is an effective disinfectant.

In order to demonstrate effectiveness disinfectants should be tested using European Standard Test Methods. Depending on the applicable area and test used, relevant log reductions are specified and must be achieved to claim effectiveness with a test method. This means a reduction in microbial numbers must be seen when compared to the number of organisms at the start of the test or, for surface tests, to a water control performed at the same time. As the numbers are large it is generally accepted that they are expressed as a logarithm. The reduction can be written as either a log value or a percentage i.e. a 5 log reduction is equivalent to a 99.999% reduction, a 3 log reduction is equivalent to 99.9% reduction.

Bacteria are microscopic free living single celled organisms. A surface contaminated with raw meat for example could have millions of bacteria per square centimetre e.g. a surface with 1,000,000 bacteria treated with a product that kills 99.9% of bacteria would still have 1000 bacteria remaining.

**If the surface were treated with a product that kills 99.999% of bacteria only 10 bacteria would remain.**

Bacterial growth rates vary depending on the surface, type and degree of soiling, temperature and presence of water. For example, E.coli under ideal conditions multiplies every 15 minutes. If conditions are less than ideal (lowering the temperature or drying the surface) the growth rate slows down.

e.g. 1,000 bacteria would increase to 2,000 after 15 minutes, after 30 minutes it would be 4,000 and after 1 hour 16,000 and 256,000 after 2 hours, **10 bacteria would only have multiplied to 2560 in the same 2 hour period.**

The presence of bacteria does not automatically lead to infection, susceptibility and the infectious dose (number of bacteria required to cause infection) are vitally important. Some bacteria will cause an infection with less than 100 cells ingested or introduced into cuts or wounds. For this reason, it is important to reduce numbers of harmful bacteria to the lowest number possible wherever the risk of infection is high.

THE FOLLOWING FIGURES APPLY IF THE NUMBER AT THE START POINT WAS 1,000,000

LOG REDUCTION	NUMBER REMAINING	PERCENTAGE REDUCTION
1	100,000	90%
2	10,000	99%
3	1,000	99.9%
4	100	99.99%
5	10	99.999%