



Evans Vanodine International plc
GLOBAL HYGIENE SOLUTIONS

SAFEZONE



MICROBIOLOGICAL PROFILE

SAFEZONE MICROBIOLOGICAL PROFILE

INTRODUCTION

SAFEZONE is an unperfumed liquid disinfectant and multi-purpose cleaner.

SAFEZONE is available as a concentrate product and in a ready-to-use (RTU) solution. The results reported in this profile have been carried out on dilutions of the concentrated product.

SAFEZONE has been tested using European Standard methods to meet specific classification/regulatory demands.

European Standard test methods EN 1276 and EN 13624 were performed in the UKAS accredited Microbiology Laboratory (Testing No. 1108) of Evans Vanodine International Plc.

Tests with additional organisms *Campylobacter jejunii* and *Listeria monocytogenes* were performed by an independent UKAS accredited laboratory.

EN 1276 uses four reference bacteria, *Enterococcus hirae*, *Escherichia coli* (*E.coli*), *Pseudomonas aeruginosa* and *Staphylococcus aureus* as representatives of the main bacterial types.

Pseudomonas aeruginosa is considered to be one of the most resistant bacteria to disinfectants and therefore the effective dilutions against this bacterium are normally used to determine recommended in-use dilutions.

PLEASE REFER TO PRODUCT LABEL FOR HOW TO USE AND FOR ALL RECOMMENDED DILUTION RATES.

CONTENTS

PAGE

BACTERICIDAL ACTIVITY IN SUSPENSION

2 - 3

Campylobacter jejunii

Enterococcus hirae

Escherichia coli

Escherichia coli "0157"

Listeria monocytogenes

Methicillin resistant *Staphylococcus aureus*

Pseudomonas aeruginosa

Salmonella pullorum

Salmonella typhimurium

Shigella sonnei

Staphylococcus aureus

YEASTICIDAL ACTIVITY IN SUSPENSION

Candida albicans

4

A glossary of microbiological and chemical terms is available on request

SAFEZONE MICROBIOLOGICAL PROFILE

Activity against bacteria in suspension using

EN 1276

BACTERIA	DISEASE / INFECTION	Bactericidal dilutions under simulated "dirty conditions"*	
		CONTACT TIMES	
		30 seconds	5 minutes
<i>Enterococcus hirae</i>	Urinary tract infections	1:200	1:400
<i>Escherichia coli</i>	Food poisoning	1:25	1:50
<i>Pseudomonas aeruginosa</i>	Opportunistic pathogen, wound, burn infections	1:25	1:25
<i>Staphylococcus aureus</i>	Skin, bone and wound infections	1:50	1:200
<i>Campylobacter jejunii</i>	Food poisoning		1:200
<i>Escherichia coli</i> "0157"	Food poisoning		1:50
<i>Listeria monocytogenes</i>	Food poisoning		1:200
<i>Methicillin resistant Staphylococcus aureus</i>	Skin, bone and wound infections		1:100
<i>Salmonella pullorum</i>	Food poisoning		1:50
<i>Salmonella typhimurium</i>	Food poisoning		1:25
<i>Shigella sonnei</i>	Dysentery		1:50

*As defined in EN 1276

SAFEZONE MICROBIOLOGICAL PROFILE

Activity against bacteria in suspension using EN 1276

BACTERIA	DISEASE / INFECTION	Bactericidal dilutions under simulated "clean conditions"*
		CONTACT TIME
		5 minutes
<i>Enterococcus hirae</i>	Urinary tract infections	1:200
<i>Escherichia coli</i>	Food poisoning	1:100
<i>Pseudomonas aeruginosa</i>	Opportunistic pathogen, wound, burn infections	1:25
<i>Staphylococcus aureus</i>	Skin, bone and wound infections	1:200

* As defined in EN 1276

TEST METHOD REFERENCE

EN 1276

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic, and institutional areas

Designed to test bactericidal products specifically for use in the Food and Catering Industry. It is carried out under "dirty" (representative of surfaces which are known to or may contain organic and/or inorganic materials) and "clean" (representative of surfaces which have received a satisfactory cleaning programme and/or are known to contain minimal levels of organic and/or inorganic materials) conditions.

Additional contact times were used as well as the obligatory test conditions.

Test Parameters: 5 minute contact time and 30 seconds, 20°C, hard water, dirty and clean conditions.

Bactericidal Criteria: ≥ 5 log reduction \equiv 99.999% reduction.

SAFEZONE MICROBIOLOGICAL PROFILE

Activity against yeast in suspension using EN 13624

BACTERIA	DISEASE / INFECTION	Bactericidal dilutions under simulated "clean conditions"*
		CONTACT TIME
		5 minutes
<i>Candida albicans</i>	Thrush	1:50

* As defined in EN 13624

TEST METHOD REFERENCE

EN 13624

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of fungicidal or yeasticidal activity in the medical area.

Designed to test fungicidal and yeasticidal products used in hospitals, health centres, dental practices, school/nursery clinics and nursing homes where disinfection or antiseptics is medically indicated.

Test Parameters: 5 minute contact time 20°C, hard water, clean conditions.
Yeasticidal Criteria: ≥ 4 log reduction \equiv 99.99% reduction.